

Locally conformal symplectic groupoids

David Iglesias-Ponte and J.C. Marrero

Departamento de Matemática Fundamental, Universidad de La Laguna
emails: diglesia@ull.es, jcmarrer@ull.es

Abstract

Locally conformal symplectic (l.c.s.) groupoids are introduced as a generalization of symplectic groupoids. We obtain some examples and we prove that l.c.s. groupoids are examples of Jacobi groupoids in the sense of [5]. Finally, we describe the Lie algebroid of a l.c.s. groupoid.

Key words: Lie groupoids, Lie algebroids, symplectic and contact groupoids, Poisson and Jacobi groupoids, Jacobi manifolds, locally conformal symplectic manifolds.

MSC 2000: 17B66, 22A22, 53D10, 53D17.

1 Introduction

A symplectic groupoid is a Lie groupoid $G \rightrightarrows M$ endowed with a symplectic structure Ω for which the graph of the partial multiplication is a lagrangian submanifold in the symplectic manifold $(G \times G \times G, \Omega \oplus \Omega \oplus -\Omega)$. This interesting class of groupoids, which was introduced in [1], arises in the integration of arbitrary Poisson manifolds. In fact, if $(G \rightrightarrows M, \Omega)$ is a symplectic groupoid then there exists a Poisson structure Λ_0 on M and the Lie algebroid AG is isomorphic to the cotangent Lie algebroid T^*M .

An interesting generalization of symplectic groupoids, as well as of Drinfeld's Poisson-Lie groups, are Poisson groupoids [8].

On the other hand, a non-degenerate 2-form on a manifold M of even dimension is said to be locally conformal symplectic (l.c.s.) if it is conformally related with a symplectic structure in some neighbourhood of every point of M (see [3, 7]). L.c.s. manifolds are interesting examples of Jacobi manifolds [3] and, in addition, they play an important role in the study of some dynamical systems, particularly conformally Hamiltonian systems (see [9]). The aim of this paper is to introduce the notion of a l.c.s. groupoid and to study some of its properties.

The paper is organized as follows. In Section 2, we recall the definition of a l.c.s. manifold and its relation with Jacobi manifolds. In Section 3, we prove that if G is a contact groupoid over a manifold M then it is possible

to introduce a l.c.s. structure and a Lie groupoid structure on $G \times \mathbb{R}$ (whose space of identity elements is $M \times \mathbb{R}$) and these structures are compatible in a certain way. These results motivate the definition of a l.c.s. groupoid in Section 4. Symplectic groupoids are l.c.s. groupoids and, furthermore, we prove, in Section 4, that a l.c.s. groupoid is a particular example of the so-called Jacobi groupoids, which were first introduced in [5] as a generalization of Poisson groupoids. Finally, in Section 5, we describe the Lie algebroid of a l.c.s. groupoid.

In this paper, we will use the definitions, notation and conventions introduced in our previous paper [5] (see Sections 1 and 2 in [5]).

2 Locally conformal symplectic manifolds

A manifold M is said to be *locally (globally) conformal symplectic (l.(g.)c.s.) manifold* if it admits a non-degenerate 2-form Ω and a closed (exact) 1-form ω such that

$$\delta\Omega = \omega \wedge \Omega. \quad (1)$$

The 1-form ω is called the *Lee 1-form* of M . It is obvious that the l.c.s. manifolds with Lee 1-form identically zero are just the symplectic manifolds (see, for example, [3, 7]).

L.c.s. manifolds are examples of Jacobi manifolds, i.e., if (M, Ω, ω) is a l.c.s. manifold then there exists a 2-vector Λ and a vector field E on M such that $[\Lambda, \Lambda] = 2E \wedge \Lambda$ and $[E, \Lambda] = 0$. In fact, the Jacobi structure (Λ, E) is given by

$$\Lambda(\alpha, \beta) = \Omega(\flat^{-1}(\alpha), \flat^{-1}(\beta)), \quad E = \flat^{-1}(\omega), \quad (2)$$

for all $\alpha, \beta \in \Omega^1(M)$, where $\flat : \mathfrak{X}(M) \longrightarrow \Omega^1(M)$ is the isomorphism of $C^\infty(M, \mathbb{R})$ -modules defined by $\flat(X) = i(X)\Omega$ (see [3]).

3 An example

Let $(G \rightrightarrows M, \eta, \sigma)$ be a *contact groupoid* over a manifold M , that is, $G \rightrightarrows M$ is a Lie groupoid over M with structural functions α, β, m and $\epsilon, \eta \in \Omega^1(G)$ is a contact 1-form on G and $\sigma : G \rightarrow \mathbb{R}$ is an arbitrary function such that if $\oplus_{TG}$ is the partial multiplication in the tangent Lie groupoid $TG \rightrightarrows TM$, then

$$\eta_{(gh)}(X_g \oplus_{TG} Y_h) = \eta_{(g)}(X_g) + e^{\sigma(g)} \eta_{(h)}(Y_h), \quad (3)$$

for $(g, h) \in G^{(2)}$ and $(X_g, Y_h) \in T_{(g,h)}G^{(2)}$ (see [2, 6]).

If $(G \rightrightarrows M, \eta, \sigma)$ is a contact groupoid then, using the associativity of $\oplus_{TG}$, we deduce that $\sigma : G \rightarrow \mathbb{R}$ is a multiplicative function, that is, $\sigma(gh) = \sigma(g) + \sigma(h)$, for $(g, h) \in G^{(2)}$. This implies that

$$\sigma \circ \epsilon = 0. \quad (4)$$

Using (3), we also deduce that (see [2])

$$\xi(\sigma) = 0, \quad (5)$$

$$\begin{aligned} (\delta\eta)_{gh}(X_g \oplus_{TG} Y_h, X'_g \oplus_{TG} Y'_h) &= (\delta\eta)_g(X_g, X'_g) + e^{\sigma(g)}(\delta\eta)_h(Y_h, Y'_h) \\ &\quad + e^{\sigma(g)}(X_g(\sigma)\eta_h(Y'_h) - X'_g(\sigma)\eta_h(Y_h)), \end{aligned} \quad (6)$$

for $(X_g, Y_h), (X'_g, Y'_h) \in T_{(g,h)}G^{(2)}$, ξ being the Reeb vector field of the contact structure η .

Now, we will obtain a l.c.s structure and a Lie groupoid structure on the manifold $G \times \mathbb{R}$, both structures compatible in a certain way.

First of all, take the Lie groupoid $G \rightrightarrows M$ and the multiplicative function $\sigma : G \rightarrow \mathbb{R}$. Then, using the multiplicative character of σ , we can define a right action of $G \rightrightarrows M$ on the canonical projection $\pi_1 : M \times \mathbb{R} \rightarrow M$ as follows

$$(x, t) \cdot g = (\alpha(g), \sigma(g) + t) \quad (7)$$

for $(x, t) \in M \times \mathbb{R}$ and $g \in G$ such that $\beta(g) = x$. Thus, we have the corresponding action groupoid $(M \times \mathbb{R}) * G \rightrightarrows M \times \mathbb{R}$ over $M \times \mathbb{R}$, where

$$(M \times \mathbb{R}) * G = \{((x, t), g) \in (M \times \mathbb{R}) \times G / \beta(g) = x\}$$

(see [4] for the general definition of an action Lie groupoid; see also [5]). Moreover, it is not difficult to prove that $(M \times \mathbb{R}) * G$ may be identified with the product manifold $G \times \mathbb{R}$ and, under this identification, the structural functions of the Lie groupoid are given by

$$\begin{aligned} \alpha_\sigma(g, t) &= (\alpha(g), \sigma(g) + t), & \text{for } (g, t) \in G \times \mathbb{R}, \\ \beta_\sigma(h, s) &= (\beta(h), s), & \text{for } (h, s) \in G \times \mathbb{R}, \\ m_\sigma((g, t), (h, s)) &= (gh, t), & \text{if } \alpha_\sigma(g, t) = \beta_\sigma(h, s), \\ \epsilon_\sigma(x, t) &= (\epsilon(x), t), & \text{for } (x, t) \in M \times \mathbb{R}. \end{aligned} \quad (8)$$

From (8) and the definition of the tangent groupoid (see, for instance, [5]), it follows that the projections $(\alpha_\sigma)^T, (\beta_\sigma)^T$, the inclusion $(\epsilon_\sigma)^T$ and the partial multiplication $\oplus_{T(G \times \mathbb{R})}$ of the tangent groupoid $T(G \times \mathbb{R}) \rightrightarrows T(M \times \mathbb{R})$ are given by

$$\begin{aligned} (\alpha_\sigma)^T(X_g + \lambda \frac{\partial}{\partial t}|_t) &= \alpha^T(X_g) + (\lambda + X_g(\sigma)) \frac{\partial}{\partial t}|_{t+\sigma(g)}, \\ (\beta_\sigma)^T(Y_h + \mu \frac{\partial}{\partial t}|_s) &= \beta^T(Y_h) + \mu \frac{\partial}{\partial t}|_s, \\ (X_g + \lambda \frac{\partial}{\partial t}|_t) \oplus_{T(G \times \mathbb{R})} (Y_h + \mu \frac{\partial}{\partial t}|_s) &= X_g \oplus_{TG} Y_h + \lambda \frac{\partial}{\partial t}|_t, \\ (\epsilon_\sigma)^T(X_x + \lambda \frac{\partial}{\partial t}|_t) &= \epsilon^T(X_x) + \lambda \frac{\partial}{\partial t}|_t, \end{aligned} \quad (9)$$

for $X_g + \lambda \frac{\partial}{\partial t}|_t \in T_{(g,t)}(G \times \mathbb{R})$, $Y_h + \mu \frac{\partial}{\partial t}|_s \in T_{(h,s)}(G \times \mathbb{R})$ and $X_x + \lambda \frac{\partial}{\partial t}|_t \in T_{(x,t)}(M \times \mathbb{R})$ (see Section 5.3.1 in [5]).

On the other hand, if $A(G \times \mathbb{R})$ is the Lie algebroid of the Lie groupoid $G \times \mathbb{R} \rightrightarrows M \times \mathbb{R}$, then using (8) and the definition of the cotangent groupoid (see [1, 5]), we deduce that the projections $\widetilde{\alpha}_\sigma, \widetilde{\beta}_\sigma$, the inclusion $\widetilde{\epsilon}_\sigma$ and the partial multiplication $\oplus_{T^*(G \times \mathbb{R})}$ in the cotangent groupoid $T^*(G \times \mathbb{R}) \rightrightarrows A^*(G \times \mathbb{R})$ are defined by

$$\begin{aligned} \widetilde{\alpha}_\sigma(\mu_g + \gamma \delta t|_t) &= (\tilde{\alpha}(\mu_g), \sigma(g) + t), \\ \widetilde{\beta}_\sigma(\nu_h + \zeta \delta t|_s) &= (\tilde{\beta}(\nu_h) - \zeta(\delta \sigma)_{\epsilon(\beta(g))}, s), \\ (\mu_g + \gamma \delta t|_t) \oplus_{T^*(G \times \mathbb{R})} (\nu_h + \zeta \delta t|_s) &= (\mu_g + \zeta(\delta \sigma)_g) \oplus_{T^*G} \nu_h + (\gamma + \zeta) \delta t|_t \\ \widetilde{\epsilon}_\sigma(\mu_x, t) &= \tilde{\epsilon}(\mu_x) + 0 \delta t|_t, \end{aligned} \tag{10}$$

for $\mu_g + \gamma \delta t|_t \in T_{(g,t)}^*(G \times \mathbb{R})$, $\nu_h + \zeta \delta t|_s \in T_{(h,s)}^*(G \times \mathbb{R})$ and $(\mu_x, t) \in A_{(x,t)}^*(G \times \mathbb{R}) \cong A_x^*G \times \mathbb{R}$, where $\tilde{\alpha}, \tilde{\beta}, \oplus_{T^*G}$ and $\tilde{\epsilon}$ are the structural functions of the cotangent Lie groupoid $T^*G \rightrightarrows A^*G$ (see Section 5.3.1 in [5]).

Next, we define on $G \times \mathbb{R}$ the 2-form Ω and the 1-form ω given by

$$\Omega = -(\pi_1^*(\delta \eta) + \pi_2^*(\delta t) \wedge \pi_1^*(\eta)), \quad \omega = -\pi_2^*(\delta t), \tag{11}$$

where $\pi_1 : G \times \mathbb{R} \rightarrow G$ and $\pi_2 : G \times \mathbb{R} \rightarrow \mathbb{R}$ are the canonical projections. Thus, we have that $(G \times \mathbb{R}, \Omega, \omega)$ is a locally conformal symplectic manifold of the first kind in the sense of [7]. Note that if (Λ, E) is the Jacobi structure on $G \times \mathbb{R}$ associated with the l.c.s. structure (Ω, ω) then

$$E = -\xi. \tag{12}$$

Now, using (3)-(6) and (8)-(12), we prove the following result.

Proposition 3.1 *Let $(G \rightrightarrows M, \eta, \sigma)$ be a contact groupoid. If $G \times \mathbb{R} \rightrightarrows M \times \mathbb{R}$ is the Lie groupoid with structural functions given by (8), the pair (Ω, ω) is defined by (11), $\bar{\sigma} = \sigma \circ \pi_1$ is the pull-back of the multiplicative function σ by the canonical projection $\pi_1 : G \times \mathbb{R} \rightarrow G$ and θ is the 1-form on $G \times \mathbb{R}$ defined by $\theta = e^{\bar{\sigma}}(\delta \bar{\sigma} - \omega)$, then we have:*

- i) $m_\sigma^* \Omega = \pi_1^* \Omega + e^{(\bar{\sigma} \circ \pi_1)} \pi_2^* \Omega;$
- ii) $\widetilde{\alpha}_\sigma \circ \omega = 0, \widetilde{\beta}_\sigma \circ \theta = 0;$
- iii) $m_\sigma^* \omega = \pi_1^* \omega, m_\sigma^* \theta = e^{(\bar{\sigma} \circ \pi_1)} \pi_2^* \theta;$
- iv) $\Lambda(\omega, \theta) = 0, (\theta + \omega - \widetilde{\epsilon}_\sigma \circ \widetilde{\beta}_\sigma \circ \omega) \circ \epsilon = 0;$

where $\pi_i : G^{(2)} \rightarrow G$, $i = 1, 2$, are the canonical projections and (Λ, E) is the Jacobi structure on $G \times \mathbb{R}$ associated with the l.c.s. structure (Ω, ω) .

4 Locally conformal symplectic and Jacobi groupoids

Motivated by Proposition 3.1, we introduce the following definition.

Definition 4.1 *Let $G \rightrightarrows M$ be a Lie groupoid with structural functions α , β , m and ϵ , (Ω, ω) be a l.c.s. structure on G , $\sigma : G \rightarrow \mathbb{R}$ be a multiplicative function and θ be the 1-form on G defined by*

$$\theta = e^\sigma(\delta\sigma - \omega). \quad (13)$$

Then, $(G \rightrightarrows M, \Omega, \omega, \sigma)$ is a l.c.s. groupoid if the following properties hold:

$$m^*\Omega = \pi_1^*\Omega + e^{(\sigma \circ \pi_1)}\pi_2^*\Omega; \quad (14)$$

$$\tilde{\alpha} \circ \omega = 0, \quad \tilde{\beta} \circ \theta = 0; \quad (15)$$

$$m^*\omega = \pi_1^*\omega, \quad m^*\theta = e^{(\sigma \circ \pi_1)}\pi_2^*\theta; \quad (16)$$

$$\Lambda(\omega, \theta) = 0, \quad (\theta + \omega - \tilde{\epsilon} \circ \tilde{\beta} \circ \omega) \circ \epsilon = 0; \quad (17)$$

where $\pi_i : G^{(2)} \rightarrow G$, $i = 1, 2$, are the canonical projections, (Λ, E) is the Jacobi structure associated with the l.c.s. structure (Ω, ω) and $\tilde{\alpha}$, $\tilde{\beta}$, $\oplus_{T^*G}$ and $\tilde{\epsilon}$ are the structural functions of the cotangent groupoid $T^*G \rightrightarrows A^*G$.

Examples 4.2 *i) If $(G \rightrightarrows M, \eta, \sigma)$ is a contact groupoid then, by Proposition 3.1, $(G \times \mathbb{R} \rightrightarrows M \times \mathbb{R}, \Omega, \omega, \bar{\sigma})$ is a l.c.s. groupoid, where the structural functions of $G \times \mathbb{R} \rightrightarrows M \times \mathbb{R}$ are defined by (8), the pair (Ω, ω) is given by (11) and the multiplicative function $\bar{\sigma}$ is $\bar{\sigma} = \sigma \circ \bar{\pi}_1$, $\bar{\pi}_1 : G \times \mathbb{R} \rightarrow G$ being the canonical projection.*

*ii) A Lie groupoid $G \rightrightarrows M$ is said to be *symplectic* if G admits a symplectic 2-form Ω in such a way that the graph of the partial multiplication in G is a Lagrangian submanifold of the symplectic manifold $(G \times G \times G, \Omega \oplus \Omega \oplus (-\Omega))$ (see [1]). This is equivalent to say that Ω satisfies the condition $m^*\Omega = \pi_1^*\Omega + \pi_2^*\Omega$. Therefore, we conclude that $(G \rightrightarrows M, \Omega)$ is a symplectic groupoid if and only if $(G \rightrightarrows M, \Omega, 0, 0)$ is a l.c.s. groupoid.*

Next, we will give the relation between l.c.s. groupoids and Jacobi groupoids. First, we will recall the definition of a Jacobi groupoid.

Let $G \rightrightarrows M$ be a Lie groupoid and $\sigma : G \rightarrow \mathbb{R}$ be a multiplicative function. Then, $TG \times \mathbb{R}$ is a Lie groupoid over $TM \times \mathbb{R}$ with structural functions given by (see Section 3 in [5])

$$\begin{aligned} (\alpha^T)_\sigma(X_g, \lambda) &= (\alpha^T(X_g), X_g(\sigma) + \lambda), \text{ for } (X_g, \lambda) \in T_g G \times \mathbb{R}, \\ (\beta^T)_\sigma(Y_h, \mu) &= (\beta^T(Y_h), \mu), \text{ for } (Y_h, \mu) \in T_h G \times \mathbb{R}, \\ (X_g, \lambda) \oplus_{TG \times \mathbb{R}} (Y_h, \mu) &= (X_g \oplus_{TG} Y_h, \lambda), \text{ if } (\alpha^T)_\sigma(X_g, \lambda) = (\beta^T)_\sigma(Y_h, \mu), \\ (\epsilon^T)_\sigma(X_x, \lambda) &= (\epsilon^T(X_x), \lambda), \text{ for } (X_x, \lambda) \in T_x M \times \mathbb{R}. \end{aligned} \quad (18)$$

On the other hand, if AG is the Lie algebroid of G then $T^*G \times \mathbb{R}$ is a Lie groupoid over A^*G with structural functions defined by (see Section 3 in [5])

$$\begin{aligned} \tilde{\alpha}_\sigma(\mu_g, \gamma) &= e^{-\sigma(g)} \tilde{\alpha}(\mu_g), \text{ for } (\mu_g, \gamma) \in T_g^*G \times \mathbb{R}, \\ \tilde{\beta}_\sigma(\nu_h, \zeta) &= \tilde{\beta}(\nu_h) - \zeta (\delta\sigma)_{\epsilon(\beta(h))|_{A_{\beta(h)}G}}, \text{ for } (\nu_h, \zeta) \in T_h^*G \times \mathbb{R}, \\ (\mu_g, \gamma) \oplus_{T^*G \times \mathbb{R}} (\nu_h, \zeta) &= ((\mu_g + e^{\sigma(g)} \zeta (\delta\sigma)_g) \oplus_{T^*G} (e^{\sigma(g)} \nu_h), \gamma + e^{\sigma(g)} \zeta) \\ \tilde{\epsilon}_\sigma(\mu_x) &= (\tilde{\epsilon}(\mu_x), 0), \text{ for } \mu_x \in A_x^*G. \end{aligned} \tag{19}$$

Definition 4.3 [5] *Let $G \rightrightarrows M$ be a Lie groupoid, (Λ, E) be a Jacobi structure on G and $\sigma : G \rightarrow \mathbb{R}$ be a multiplicative function. Then, $(G \rightrightarrows M, \Lambda, E, \sigma)$ is a Jacobi groupoid if the homomorphism $\#_{(\Lambda, E)} : T^*G \times \mathbb{R} \rightarrow TG \times \mathbb{R}$ given by*

$$\#_{(\Lambda, E)}(\mu_g, \gamma) = (\#_\Lambda(\mu_g) + \gamma E_g, -\mu_g(E_g))$$

*is a morphism of Lie groupoids over some map $\varphi_0 : A^*G \rightarrow TM \times \mathbb{R}$, where the structural functions of the Lie groupoid structure on $T^*G \times \mathbb{R} \rightrightarrows A^*G$ (respectively, $TG \times \mathbb{R} \rightrightarrows TM \times \mathbb{R}$) are given by (19) (respectively, (18)).*

Remark 4.4 A Poisson groupoid is a Jacobi groupoid $(G \rightrightarrows M, \Lambda, E, \sigma)$ with $E = 0$ and $\sigma = 0$ (see [5]).

A characterization of a Jacobi groupoid is the following one. If $G \rightrightarrows M$ is a Lie groupoid and $\sigma : G \rightarrow \mathbb{R}$ is a multiplicative function then T^*G is a Lie groupoid over A^*G with structural functions given by

$$\begin{aligned} \tilde{\alpha}_\sigma^*(\mu_g) &= e^{-\sigma(g)} \tilde{\alpha}(\mu_g), \text{ for } \mu_g \in T_g^*G, \\ \tilde{\beta}_\sigma^*(\nu_h) &= \tilde{\beta}(\nu_h), \text{ for } \nu_h \in T_h^*G, \\ \mu_g \oplus_{T^*G}^\sigma \nu_h &= \mu_g \oplus_{T^*G} e^{\sigma(g)} \nu_h, \text{ if } \tilde{\alpha}_\sigma^*(\mu_g) = \tilde{\beta}_\sigma^*(\nu_h), \\ \tilde{\epsilon}_\sigma^*(\mu_x) &= \tilde{\epsilon}(\mu_x), \text{ for } \mu_x \in A_x^*G. \end{aligned} \tag{20}$$

We call this Lie groupoid the σ -cotangent groupoid. Note that the canonical inclusion $T^*G \rightarrow T^*G \times \mathbb{R}$, $\mu_g \mapsto (\mu_g, 0)$, is a monomorphism of Lie groupoids.

Proposition 4.5 *Let $G \rightrightarrows M$ be a Lie groupoid, (Λ, E) be a Jacobi structure on G and $\sigma : G \rightarrow \mathbb{R}$ be a multiplicative function. Then, $(G \rightrightarrows M, \Lambda, E, \sigma)$ is a Jacobi groupoid if and only if the following conditions hold:*

- i) $\#_\Lambda : T^*G \rightarrow TG$ is a Lie groupoid morphism over some map $\tilde{\varphi}_0 : A^*G \rightarrow TM$ from the σ -cotangent groupoid $T^*G \rightrightarrows A^*G$ to the tangent Lie groupoid $TG \rightrightarrows TM$.

ii) E is a right-invariant vector field on G and $E(\sigma) = 0$.

iii) If $X_0 \in \Gamma(AG)$ is the section of the Lie algebroid AG satisfying $E = -\overrightarrow{X_0}$, we have that

$$\#_\Lambda(\delta\sigma) = \overrightarrow{X_0} - e^{-\sigma}\overleftarrow{X_0}.$$

Proof: Suppose that $(G \rightrightarrows M, \Lambda, E, \sigma)$ is a Jacobi groupoid. Then, proceeding as in the proof of Proposition 4.4 in [5], we deduce that i), ii) and iii) hold.

A similar computation proves the converse. $\square$

Now, we will show that a l.c.s. symplectic groupoid is a particular example of a Jacobi groupoid.

Theorem 4.6 *Let $G \rightrightarrows M$ be a Lie groupoid, (Ω, ω) be a l.c.s. structure on G and $\sigma : G \rightarrow \mathbb{R}$ be a multiplicative function. If (Λ, E) is the Jacobi structure associated with the l.c.s. structure (Ω, ω) then $(G \rightrightarrows M, \Omega, \omega, \sigma)$ is a l.c.s. groupoid if and only if $(G \rightrightarrows M, \Lambda, E, \sigma)$ is a Jacobi groupoid.*

Proof: Assume that $(G \rightrightarrows M, \Omega, \omega, \sigma)$ is a l.c.s. groupoid. If $\mu_g \in T_g^*G$ and $\nu_h \in T_h^*G$ satisfy the relation $\tilde{\alpha}_\sigma^*(\mu_g) = \tilde{\beta}_\sigma^*(\nu_h)$, then using (2), (14) and the definition of the partial multiplication $\oplus_{TG}$ in the tangent Lie groupoid $TG \rightrightarrows TM$, we obtain that $\alpha^T(\#_\Lambda(\mu_g)) = \beta^T(\#_\Lambda(\nu_h))$ and, in addition,

$$\begin{aligned} (i(\#_\Lambda(\mu_g \oplus_{T^*G}^\sigma \nu_h))\Omega_{(gh)})(X_g \oplus_{TG} Y_h) \\ = (i(\#_\Lambda(\mu_g) \oplus_{TG} \#_\Lambda(\nu_h))\Omega_{(gh)})(X_g \oplus_{TG} Y_h), \end{aligned}$$

for $(X_g, Y_h) \in T_{(g,h)}G^{(2)}$. Thus (see (2)), it follows that $\#_\Lambda(\mu_g \oplus_{T^*G}^\sigma \nu_h) = \#_\Lambda(\mu_g) \oplus_{TG} \#_\Lambda(\nu_h)$ and, therefore, the map $\#_\Lambda : T^*G \rightarrow TG$ is a Lie groupoid morphism over some map $\tilde{\varphi}_0 : A^*G \rightarrow TM$, between the σ -cotangent groupoid $T^*G \rightrightarrows A^*G$ and the tangent groupoid $TG \rightrightarrows TM$. In particular, this implies that

$$\alpha^T \circ \#_\Lambda = \tilde{\varphi}_0 \circ \tilde{\alpha}_\sigma^*, \quad \beta^T \circ \#_\Lambda = \tilde{\varphi}_0 \circ \tilde{\beta}_\sigma^*. \quad (21)$$

Now, from (2), we deduce that $E = -\#_\Lambda(\omega)$. Using this relation, (15) and (21), we have that the vector field E is α -vertical

Next, suppose that $(g, h) \in G^{(2)}$ and denote by $R_h : G_{\beta(h)} \rightarrow G_{\alpha(h)}$ the right-translation by h . Then, (2) and (16) imply that

$$(i(E_{(gh)})\Omega_{(gh)})(X_g \oplus_{TG} Y_h) = (i((R_h)_*^g(E_{(g)}))\Omega_{(gh)})(X_g \oplus_{TG} Y_h),$$

for $(X_g, Y_h) \in T_{(g,h)}G^{(2)}$. Consequently, E is a right-invariant vector field and there exists $X_0 \in \Gamma(AG)$ such that $E = -\overrightarrow{X_0}$.

On the other hand, if X_{e^σ} is the hamiltonian vector field of the function e^σ , $X_{e^\sigma} = e^\sigma \#_\Lambda(\delta\sigma) + e^\sigma E$, it is clear that $X_{e^\sigma} = \#_\Lambda(\theta)$. Using this equality,

(15), (16), (21) and proceeding as in the proof of the fact that E is right-invariant, we conclude that X_{e^σ} is a left-invariant vector field. Moreover, if x is a point of M , then relation (17) implies that $X_{e^\sigma}(\epsilon(x)) = -\overleftarrow{X}_0(\epsilon(x))$. Thus, $\#_\Lambda(\delta\sigma) = \overrightarrow{X}_0 - e^{-\sigma}\overleftarrow{X}_0$.

Finally, since $\Lambda(\omega, \theta) = 0$ and $E = -\#_\Lambda(\omega)$, we obtain that $E(\sigma) = 0$. Therefore, $(G \rightrightarrows M, \Lambda, E, \sigma)$ is a Jacobi groupoid.

In a similar way, we prove the converse. $\square$

Remark 4.7 Using Theorem 4.6 we directly deduce that a symplectic groupoid is a Poisson groupoid. This result was proved in [8].

5 The Lie algebroid of a l.c.s. groupoid

Let $(G \rightrightarrows M, \Omega, \omega, \sigma)$ be a l.c.s. groupoid and θ the 1-form on G given by (13). Then, the 1-form $e^{-\sigma}\theta$ is closed and since $\tilde{\beta} \circ \theta = 0$, it follows that θ is basic with respect to the projection α . Thus, there exists a unique 1-form θ_0 on M such that $\alpha^*\theta_0 = e^{-\sigma}\theta$. It is clear that θ_0 is closed.

Now, denote by (Λ, E) the Jacobi structure on G associated with the l.c.s. structure (Ω, ω) . Then, $\#_\Lambda(\theta)$ is the hamiltonian vector field X_{e^σ} of the function e^σ . Moreover, from Theorem 4.6, and using the results in [5] (see Proposition 5.6 in [5]), we deduce that there exists a Jacobi structure (Λ_0, E_0) on M in such a way that the couple (α, e^σ) is a conformal Jacobi morphism between the Jacobi manifolds (G, Λ, E) and (M, Λ_0, E_0) . This implies that

$$\begin{aligned} \#_{\Lambda_0(\alpha(g))}(\theta_0(\alpha(g))) &= e^{\sigma(g)}(\alpha_g^T \circ \#_{\Lambda(g)} \circ (\alpha_g^T)^*)(\theta_0(\alpha(g))) \\ &= \alpha_g^T(X_{e^\sigma}(g)) = E_0(\alpha(g)), \end{aligned}$$

for $g \in G$, where $(\alpha_g^T)^* : T_{\alpha(g)}^*M \rightarrow T_g^*G$ is the adjoint map of the tangent map $\alpha_g^T : T_gG \rightarrow T_{\alpha(g)}M$. Therefore, we have proved the following result.

Proposition 5.1 *Let $(G \rightrightarrows M, \Omega, \omega, \sigma)$ be a l.c.s. groupoid and θ the 1-form on G given by (13). Then, there exists a unique 1-form θ_0 on M such that $\alpha^*\theta_0 = e^{-\sigma}\theta$. Furthermore, θ_0 is closed and $\#_{\Lambda_0}(\theta_0) = E_0$.*

Next, we will describe the Lie algebroid associated with a l.c.s. groupoid.

Theorem 5.2 *Let $(G \rightrightarrows M, \Omega, \omega, \sigma)$ be a l.c.s. groupoid, AG be the Lie algebroid of G , (Λ, E) be the Jacobi structure on G associated with the l.c.s. structure (Ω, ω) and (Λ_0, E_0) be the corresponding Jacobi structure on M . Then, the map $\Psi : \Omega^1(M) \rightarrow \mathfrak{X}_L(G)$ between $\Omega^1(M)$ and the space of left-invariant*

vector fields on G defined by $\Psi(\mu) = e^\sigma \#_\Lambda(\alpha^* \mu)$ induces an isomorphism between the vector bundles T^*M and AG . Under this isomorphism, the Lie bracket on $\Gamma(AG) \cong \mathfrak{X}_L(G)$ and the anchor map of AG are given by

$$\begin{aligned} \llbracket \mu, \nu \rrbracket_{(\Lambda_0, E_0, \theta_0)} &= \mathcal{L}_{\#_{\Lambda_0}(\mu)} \nu - \mathcal{L}_{\#_{\Lambda_0}(\nu)} \mu - \delta(\Lambda_0(\mu, \nu)) \\ &\quad - i(E_0)(\mu \wedge \nu) - \Lambda_0(\mu, \nu) \theta_0, \\ \tilde{\#}_{(\Lambda_0, E_0, \theta_0)}(\mu) &= \#_{\Lambda_0}(\mu), \end{aligned}$$

for $\mu, \nu \in \Omega^1(M)$, where θ_0 is the 1-form on M considered in Proposition 5.1.

Proof: Let μ be a 1-form on M . Since the map $\#_\Lambda : T^*G \rightarrow TG$ is a morphism between the σ -cotangent groupoid and the tangent groupoid $TG \rightrightarrows TM$, we obtain that the vector field $\tilde{X} = \Psi(\mu)$ is β -vertical. Moreover, if $(g, h) \in G^{(2)}$ and $L_g : G^{\alpha(g)} \rightarrow G^{\beta(g)}$ is the left-translation by g then, using (14), we deduce that

$$(i(\tilde{X}_{(gh)})\Omega_{(gh)})(Y_g \oplus_{TG} Z_h) = (i((L_g)_*^h(\tilde{X}_{(h)}))\Omega_{(gh)})(Y_g \oplus_{TG} Z_h).$$

for $(Y_g, Z_h) \in T_{(g,h)}G^{(2)}$. This proves that $\tilde{X} \in \mathfrak{X}_L(G)$.

Conversely, if $\tilde{X} \in \mathfrak{X}_L(G)$ and $\tilde{\mu}$ is the 1-form on G defined by $\tilde{\mu} = -i(\tilde{X})\Omega$ then, from (14), it follows that $e^\sigma \alpha^* \mu = \tilde{\mu}$, μ being the 1-form on M given by $\mu = \epsilon^* \tilde{\mu}$. This implies that $\Psi(\mu) = \tilde{X}$.

On the other hand, using that the map $\#_\Lambda : \Omega^1(G) \rightarrow \mathfrak{X}(G)$ is an isomorphism of $C^\infty(G, \mathbb{R})$ -modules, we conclude that Ψ is an isomorphism of $C^\infty(M, \mathbb{R})$ -modules.

Now, suppose that $X, Y \in \Gamma(AG)$. We have that the left-invariant vector field $\tilde{X}$ is α -projectable to a vector field $a(X)$ on M . In addition, if μ and ν are the 1-forms on M satisfying $\Psi(\mu) = \tilde{X}$ and $\Psi(\nu) = \tilde{Y}$, then a long computation, using (1) and Definition 4.1, shows that

$$a(X) = \tilde{\#}_{(\Lambda_0, E_0, \theta_0)}(\mu), \quad i([\tilde{X}, \tilde{Y}])\Omega = -e^\sigma \alpha^* \llbracket \mu, \nu \rrbracket_{(\Lambda_0, E_0, \theta_0)}.$$

This ends the proof of our result. $\square$

Remark 5.3 Let $(G \rightrightarrows M, \Omega)$ be a symplectic groupoid. Then, the Jacobi structure on M is Poisson, that is, $E_0 = 0$ (see Section 5.2 in [5]) and the 1-form θ_0 on M identically vanishes. Thus (see Theorem 5.2), AG is isomorphic to the cotangent Lie algebroid T^*M . This result was proved in [1].

Acknowledgments

Research partially supported by DGICYT grant BFM 2000-0808. D. Iglesias-Ponte wishes to thank the Spanish Ministry of Education and Culture for an FPU grant.

References

- [1] A. COSTE, P. DAZORD, A. WEINSTEIN, “Groupoïdes symplectiques”, *Pub. Dép. Math. Lyon* **2/A** (1987) 1–62.
- [2] P. DAZORD, “Sur l’intégration des algèbres de Lie locales et la préquantification”, *Bull. Sci. Math.* **121** (1990) 423–462.
- [3] F. GUÉDIRA, A. LICHNEROWICZ, “Géométrie des algèbres de Lie locales de Kirillov”, *J. Math. pures et appl.* **63** (1984) 407–484.
- [4] P.J. HIGGINS, K.C.H. MACKENZIE, “Algebraic constructions in the category of Lie algebroids”, *J. Algebra* **129** (1990) 194–230.
- [5] D. IGLESIAS-PONTE, J.C. MARRERO, “Jacobi groupoids and generalized Lie bialgebroids”, *Preprint* (2002) math.DG/0208032.
- [6] Y. KERBRAT, Z. SOUICI-BENHAMMADI, “Variétés de Jacobi et groupoïdes de contact”, *C.R. Acad. Sci. Paris* **317** Sér. I (1993) 81–86.
- [7] I. VAISMAN, “Locally conformal symplectic manifolds”, *Internat. J. Math. & Math. Sci.* **8** (1985) 521–536.
- [8] A. WEINSTEIN, “Coisotropic calculus and Poisson groupoids”, *J. Math. Soc. Japan* **40** (1988) 705–727.
- [9] M.P. WOJTKOWSKI, C. LIVERANI, “Conformally symplectic dynamics and symmetry of the Lyapunov spectrum”, *Comm. Math. Phys.* **194** (1998) 47–60.